

AD_____

Award Number: W81XWH-05-1-0233

TITLE: A proteomic approach to identify phosphorylation-dependent targets of BRCT domains

PRINCIPAL INVESTIGATOR: Zhou Songyang, Ph.D.

CONTRACTING ORGANIZATION: Baylor College of Medicine
Houston, Texas 77030

REPORT DATE: March 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01/03/2006		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 1 Mar 2005 – 28 Feb 2006	
A proteomic approach to identify phosphorylation-dependent targets of BRCT domains				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-05-1-0233	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Zhou Songyang, Ph.D. E-mail: songyang@bcm.tmc.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Baylor College of Medicine Houston, Texas 77030				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT BRCA1 C-terminal (BRCT) domains are novel phosphopeptide binding modules. Cancer-associated missense and deletion mutations have been found in the BRCT repeat regions of BRCA1, suggesting an essential role of BRCT domains in regulating BRCA1 activity. In addition, BRCT domains are found in many proteins that regulate DNA damage repair, cell cycle, and genome stability, implying a more global role of BRCT domains in genome stability surveillance. These results suggest that the BRCT domain acts as a sensor to protein phosphorylation in response to DNA damage, recruits phosphorylated cellular targets, and mediates signaling complex formation. However, the identities of the in vivo BRCT domain targets are largely unknown. In order to understand the role of phosphorylation in protein-protein interactions, we developed several approaches utilizing peptide libraries and peptide arrays. We propose to use these methods to systematically identify phosphoproteins that can interact with BRCT domains. In addition to potential new regulators of genome stability, the approaches can identify phosphorylated sequences on proteins that are important for DNA damage responses and cell cycle. Such information should prove valuable, especially for the development of new screening strategies, drug targets, and treatment for breast cancer.					
15. Subject Terms (keywords previously assigned to proposal abstract or terms which apply to this award) BRCT domain, peptide library, OPAL, peptide array, proteomics, genome wide, signal transduction pathways, androgen receptor					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

Cover.....	
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusions.....	7
References.....	7

Introduction

Understanding the molecular and cellular mechanisms that trigger breast cancer is essential to the prevention and treatment of this disease. The BRCA1 C-terminal (BRCT) domain, was first identified in BRCA1 (1, 3). Cancer associated missense and deletion mutations have been found in the BRCT repeat regions of BRCA1, suggesting an important role of BRCT domains in regulating BRCA1 activity (16, 23). In addition, the BRCT domain is found in many proteins that regulate DNA damage repair, cell cycle, and genome stability, implying a more global role of BRCT domains in genome stability surveillance (1, 3). Consistent with this notion, the BRCT domain has been shown to mediate protein-protein interactions. For example, BRCT domains of BRCA1 associate with helicase BACH1 and CtBP interacting protein CTIP (4, 49). Recently, our lab and others have discovered that BRCT domains are novel phosphopeptide binding modules (20, 32, 48). BRCA1 BRCT domains associate with residue Ser990 on BACH1 in a phosphorylation-dependent manner. Furthermore, we found that several other BRCT domains including those from MDC1 and tumor suppressor BARD1 can bind specific phosphorylated peptides (12, 19, 38). These findings suggest that the BRCT domain recruits phosphorylated cellular targets and mediates signaling complex formation. However, the identities of the *in vivo* BRCT domain targets are largely unknown. In this application, we propose to use peptide libraries and peptide arrays to systematically identify phosphoproteins that can interact with BRCT domains. We may uncover potential new regulators of genome stability; more importantly, the approach can identify phosphorylated sequences on proteins that are important for DNA damage responses and cell cycle. Such information will help us to understand the mechanism of how protein phosphorylation modulates DNA damage responses and cell cycle in breast epithelial cells. In addition, it should prove invaluable for the development of new screening strategies and treatment for breast cancer.


Body

A. For Task 1, we proposed to identify phosphorylated peptide sequences that specifically bind BRCA1 and BARD1 BRCT domains. This will be achieved by (1) establishing Oriented Peptide Array Libraries (OPAL) and using OPAL to examine the specificity of BRCA1 and BARD1, and (2) by establishing peptide chips to screen for BRCA1 and BARD1 BRCT domain binding targets.

1) For the first part of Task 1, we focused on BRCT domain fusion proteins and OPAL arrays, both of which are key reagents.

1.1 First generation OPAL was synthesized on cellulose paper and worked well with SH2 domains and kinases. We therefore carried out analysis of BRCA1 and BARD1 BRCT domain fusion proteins using cellulose paper based OPAL arrays. The set up of OPAL is depicted in Figure 1a (left panel) and the results of BARD1 BRCT domain is shown in Figure 1 (right panel). Clearly, the BARD1 BRCT domain had no specific sequence preference for the amino terminus of the peptide. Furthermore, it showed preference for Asp and Glu residues at the carboxy terminus. This result is consistent with our published finding that BARD1 prefers acidic residues at position C-terminus to the phosphoserine

2) The second part of Task 1 seeks to screen for BARD1 binding peptides. To accomplish this, we synthesized an array (peptide chip) containing ~500 potential DNA damage phosphorylation sites selected from a number of proteins (Figure 3). We performed binding assays using BARD1BRCT domain and control proteins. The approach was validated by our identification of a number of target sites. For example, phosphopeptide sequences from BACH1 were identified for BRCA1 BRCT domain using this method.



There were a few reactive spots on the array when probed with BARD1 (Figure 3). However, the same spots were also positive in the control array experiments probed with non-BRCT proteins. Therefore, none of these phosphopeptides bound specifically to BARD1 BRCT domain, suggesting that they are not targets of BARD1. We are currently generating larger arrays or developing new methods to identify BARD1 BRCT targets.

B. For Task 2, we proposed to characterize breast cancer genes and

BRCT binding sites identified in Task 1, by RNAi and by determining the role of BRCT-phosphopeptide interaction in DNA damage. While we are optimizing screening conditions and generating assay reagents, we have conducted more detailed analysis of BARD1.

Three retroviral RNAi vectors for BARD1 have been made. We also have obtained two different antibodies for human BARD1. We currently analyzing these antibodies to make sure that they can both immunoprecipitate and western blot BARD1, which should help us to identify the RNAi vectors that significantly knockdown BARD1 expression.

Key Research Accomplishments

A number of key findings have resulted from the work so far:

- Analyzed BRCA1 and BARD1 BRCT domains using OPAL spotted on cellulose
- Analyzed BARD1 BRCT domain using peptide chips containing ~500 peptides
- Constructed RNAi vectors for BARD1
- Engineered Gateway pENTR and Destination vectors for epitope-tagged BRCT domain expression
- Preliminary data on BARD1 BRCT binding sites
- Demonstrated feasibility of using OPAL to screen for phospho-binding sites
- Identification of problems associated with cellulose based binding assays
- We have already obtained and verified a collection of GST-BRCT domain fusion proteins
- We showed feasibility of peptide chips containing ~500 peptides for binding analysis

Reportable Outcomes

We have utilized the OPAL and peptide chips approach to investigate the binding specificities of BRCA1 and BARD1 BRCT domains. A motif for BARD1 BRCT domain recognition was identified.

Meeting abstract/presentation

Specificity of BRCT domains and breast cancer. (2005) Maria Rodriguez, Zhou Songyang. SCBA Annual Research Conference. Houston, TX

Conclusions

In summary, we have conducted pilot screens of BRCT domain interacting sequences using peptide chips. Furthermore, we are in the process of optimizing OPAL for BRCT interaction screens. We determined that a different surface (other than cellulose) or longer linker may be necessary for our proposed screen. Lastly, we have generated the necessary reagents (such as RNAi vectors, antibodies) for the second phase of our project. The information obtained from our studies should prove especially useful for the development of new and effective screening strategies, drug targets, and treatment for breast cancer.

References

1. **Ellisen, L. W., and D. A. Haber.** 1998. Hereditary breast cancer. *Annu Rev Med* **49**:425-36.
2. **Rahman, N., and M. R. Stratton.** 1998. The genetics of breast cancer susceptibility. *Annu Rev Genet* **32**:95-121.

3. **Liu, D., Yang, X., Yang, D., Songyang, Z.** 2000. Genetic screens in mammalian cells using Enhanced Retroviral Mutagens. *Oncogene* **19**:5964-72.
4. **Yoshida K, Weichselbaum R, Kharbanda S, Kufe D.** 2000. Role for Lyn tyrosine kinase as a regulator of stress-activated protein kinase activity in response to DNA damage. *Mol Cell Biol* **20**:5370-80.
5. **Maruo A, Oishi I, Sada K, Nomi M, Kurosaki T, Minami Y, Yamamura H.** 1999. Protein tyrosine kinase Lyn mediates apoptosis induced by topoisomerase II inhibitors in DT40 cells. *Int Immunol* **11**:1371-80.
6. **Kharbanda S, Saleem A, Yuan ZM, Kraeft S, Weichselbaum R, Chen LB, Kufe D.** 1996. Nuclear signaling induced by ionizing radiation involves colocalization of the activated p56/p53lyn tyrosine kinase with p34cdc2. *Cancer Res* **56**:3617-21.
7. **Kharbanda S, Yuan ZM, Taneja N, Weichselbaum R, Kufe D.** 1994. **p56/p53lyn tyrosine kinase activation in mammalian cells treated with mitomycin C.** *Oncogene* **9**:3005-11.
8. **Wu, L. C., Z. W. Wang, J. T. Tsan, M. A. Spillman, A. Phung, X. L. Xu, M. C. Yang, L. Y. Hwang, A. M. Bowcock, and R. Baer.** 1996. Identification of a RING protein that can interact in vivo with the BRCA1 gene product. *Nat Genet* **14**:430-40.
9. **Thai, T. H., F. Du, J. T. Tsan, Y. Jin, A. Phung, M. A. Spillman, H. F. Massa, C. Y. Muller, R. Ashfaq, J. M. Mathis, D. S. Miller, B. J. Trask, R. Baer, and A. M. Bowcock.** 1998. Mutations in the BRCA1-associated RING domain (BARD1) gene in primary breast, ovarian and uterine cancers. *Hum Mol Genet* **7**:195-202.
10. **Dichtl B, Blank D, Sadowski M, Hubner W, Weiser S, Keller W.** 2002. Yhh1p/Cft1p directly links poly(A) site recognition and RNA polymerase II transcription termination. *EMBO J* **21**: 4125-35.
11. **Salinas CA, Sinclair DA, O'Hare K, Brock HW.** 1998. Characterization of a Drosophila homologue of the 160-kDa subunit of the cleavage and polyadenylation specificity factor CPSF. *Mol Gen Genet* Apr;257(6):672-80.
12. **Takagaki Y, Manley JL.** 2000. Complex protein interactions within the human polyadenylation machinery identify a novel component. *Mol Cell Biol* **20**:1515-25.